

Fields of Innovation 2021

Event

Powered by Syngenta
Vegetable Seeds

Thank you
for visiting
and enjoy your
Field Book!

The Syngenta logo features the word "syngenta" in a dark blue, lowercase sans-serif font. A small green leaf icon is positioned above the letter 'n'. A registered trademark symbol (®) is located at the end of the word.

Content of your Field Book

Bolikor

Rosberg SGW0541

Succesor SGW0359

Katator SGW0303

Terminator SGW0247

Toreador SGW0417



Powered by
Syngenta Vegetable Seeds

White Cabbage



Bolikor

- Early variety for open field production
- Around 60-65 growing days
- Nice round dark green head weight (2-2,5 kg)
- Long field standing ability
- Suitable to grow in difficult early spring conditions

Variety	Segment	Resistances
Bolikor	Fresh Early	HR: Foc 1

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

syngenta[®]

Powered by
Syngenta Vegetable Seeds

White Cabbage



Rosberg SGW0541

- Adaptability to cold spring conditions tunnel and open field
- 50-55 growing days, strong bolting tolerance
- Good field standing ability and density of heads
- Round heads weight 1,2-1,8 kg

Variety	Segment	Resistances
Rosberg SGW0541	Fresh Early	HR: Foc 1

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with its own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

syngenta[®]

Powered by
Syngenta Vegetable Seeds

White Cabbage



Successor SGW0359

- Medium fresh and short storage
- Variety in Agressor segment recommended for areas with higher pressure of Xanthomonas
- More round heads weight 3-5 kg, perfect for supermarkets
- Fresh market and processing
- Excellent performance in continental conditions

Variety	Segment	Resistances
Successor SGW0359	Autumn / Storage	HR: Foc 1, Strong against Tt

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

syngenta®

Powered by
Syngenta Vegetable Seeds

White Cabbage



Katator SGW0303

- Quisor type for summer production
- Very strong against thrips
- Head weight 2,5-3,5 kg
- Fresh market variety

Variety	Segment	Resistances
Katator SGW0303	Fresh Medium Late	HR: Foc 1, Strong against Tt

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

syngenta®

Powered by
Syngenta Vegetable Seeds

White Cabbage



Terminator SGW0247

- New medium variety for Europe
- In continental stress growing conditions
- Maturity 100-110 days
- Head size 4-5 kg Strong
- Healthy plant Strong against thrips and Xcc

Variety	Segment	Resistances
Terminator SGW0247	Fresh Medium Late	HR: Foc 1, Strong against Tt and Xcc

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

syngenta®

Powered by
Syngenta Vegetable Seeds

White Cabbage



Toreador SGW0417

- Medium fresh segment variety, 90-95 growing days
- Head weight 2-3 kg, excellent for supermarkets
- Strong against thrips, outstanding performance across different geographies
- Upright leaves and high uniformity
- Perfect for mechanical harvest

Variety	Segment	Resistances
Toreador SGW0417	Fresh Medium Late	HR: Foc 1, Strong against Tt and Xcc

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.


Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

syngenta[®]



**You can always
visit us again
anytime you want**

foi.syngentavegetables.com

syngenta®