

Fields of Innovation

September 21 - 25, 2020

Event

Powered by Syngenta
Vegetable Seeds

Thank you
for visiting
and enjoy your
Field Book!

syngenta®

Content of your Field Book

Rhodos

Balard SGD15-0086

Batory

Besty

Monflor

Aldoran SGC2105

Almagro

Andromeda SGC2074

Amidala SGC2834

Astrotek SGC2130

Cleozil SGC2101

Draconis

Electra SGC3106

Guideline

Koten SGC3089



Content of your Field Book

Space Dream SGC2095

Spacestar Gold SGC2142

Wonder SGC3096

Bogart SGK0008

Brando SGK0024

Fonda SGK0007

Hey Melis

Marcantus

Eldoret SGW0431

Pruktor Cemes

Bolikor

Rosberg SGW0541

Succesor SGW0359

Webber SGW0508

Katator SGW0303



Content of your Field Book

Terminator SGW0247

Toreador SGW0417

Kilastress SGW0377

Kepler

Storidor

Red River SGX0057

Atractum SGS0325

Coradi SGS0323

Cordesa

Corripa SGS0294

Milandina SGS0318

Manitoba



Squash



Rhodos

- Saves time to harvest as the plant is really open and erect
- High yielding variety, with good earliness, to maximize your profitability
- Excellent fruit quality, with a dark green colour, to satisfy your customers

Variety	Segment	Resistances
Rhodos	General	IR : Gc , Px ,CMV , WMV , ZYMV

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

Powered by
Syngenta Vegetable Seeds

Broccoli



Balard SGD15-0086

- Variety for Europe, China, Australia
- Dark green head colour with nice shape and firmness
- Excellent vigour and field standing ability
- Good shelf life
- Top quality for spring, autumn and winter harvest

Variety	Segment	Resistances
Balard SGD15-0086	Head	

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

syngenta[®]

Broccoli



Batory

- 75-80 growing days for autumn harvest for North Europe
- Suitable for processing industry and fresh market variety
- Broad harvest window
- Suitable for different growing conditions spring and summer
- Outstanding performance during hot 2019 season

Variety	Segment	Resistances
Batory	Head	

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

Broccoli



Besty

- 70-75 growing days, easy to grow variety for fresh market for Europe, Africa, Middle East and India
- Summer production in continental conditions
- Mushroom head shape with small beads
- In Montop segment for Africa and Middle East

Variety	Segment	Resistances
Besty	Head	

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

Powered by
Syngenta Vegetable Seeds

Broccoli



Variety
video



Monflor

- First variety from EasyBroq™
- Early multifloret broccoli
- Medium maturity, 55-60 growing days
- Very high floret quality, no yellow borders
- Flexibility in harvest with short, medium and long stems

Variety	Segment	Resistances
Monflor	Specialties	

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

syngenta®

Powered by
Syngenta Vegetable Seeds

Cauliflower



Variety
video



Highlights
video



Aldoran SGC2105



- Medium maturity variety for Europe, Africa and Middle East 70-75 growing days
- Keeps white colour even when no self protection
- No need to cover the curds, labour saving for the grower
- Strong plant and good self protection
- Recommended for spring and autumn harvest

Variety	Segment	Resistances
Aldoran SGC2105	Fresh	

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

syngenta[®]

Cauliflower



Variety
video



Almagro

- Leading variety for summer continental growing conditions in North-East Europe
- Medium maturity - 75-80 growing days
- For fresh market, harvest extension for Lecanu
- Strong plants and white curd colour
- Very good self-protection

Variety	Segment	Resistances
Almagro	Fresh	

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

Powered by
Syngenta Vegetable Seeds

Cauliflower



Variety
video



Highlights
video



Andromeda SGC2074



- Medium maturity variety 65-70 growing days
- Keeps white colour even when no self protection
- No need to cover the curds, labour saving for the grower
- Suitable for different growing conditions in EAME region

Variety	Segment	Resistances
Andromeda SGC2074	Fresh	

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

syngenta[®]

Powered by
Syngenta Vegetable Seeds

Cauliflower



Variety
video



Amidala SGC2834

- Mid cycle autumn variety for South Europe and summer period in Middle East
- Very good curd quality in hot conditions
- Reliable growing period
- Perfect self protection
- High quality curds for fresh market

Variety	Segment	Resistances
Amidala SGC2834	Fresh	

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

syngenta[®]

Powered by
Syngenta Vegetable Seeds

Cauliflower


Variety
video



Astrotek SGC2130

- Medium cycle variety 70-75 days for Europe
- Erected strong plant
- Very healthy variety adapted to different growing conditions
- Firm, white curd, well protected
- For fresh market

Variety	Segment	Resistances
Astrotek SGC2130	Fresh	

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation



Powered by
Syngenta Vegetable Seeds

Cauliflower


Variety
video



Cleozil SGC2101

- Early Clarina type for fresh market for north Europe
68-73 growing days
- Medium plant vigour
- Very high harvestable yield
- High quality curds

Variety	Segment	Resistances
Cleozil SGC2101	Fresh	HR:Pb

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation



Cauliflower



Draconis

- Variety suitable for USA
- Vigorous plant with excellent self-protection (wrap)
- Bright white dense curd with great quality
- Medium cycle cauliflower for overwinter and spring slot California
- Uniform maturity

Variety	Segment	Resistances
Draconis	Fresh	

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

Powered by
Syngenta Vegetable Seeds

Cauliflower



Variety
video



Electra SGC3106

- New variety in Lecanu segment for Europe
- Stable performance in continental growing condition
- Well balanced plant
- Heavy white curd
- Suitable for fresh and procesing

Variety	Segment	Resistances
Electra SGC3106	Fresh	

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

syngenta[®]

Cauliflower



Guideline

- Winner in Germany
- Medium cycle variety
- 70-75 growing days with perfect curd setting in summer
- Very reliable in growing days
- Very good curd quality
- For the whole season production in North Europe

Variety	Segment	Resistances
Guideline	Fresh	

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

Cauliflower



Koten SGC3089

- Late maturity variety 105-115 growing days
- For autumn harvest in France, November
- Strong plant vigour and sanitary
- Very good self protection and curd quality
- For fresh market

Variety	Segment	Resistances
Koten SGC3089	Fresh	

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

Powered by
Syngenta Vegetable Seeds

Cauliflower



Space Dream SGC2095

- Medium cycle variety for Europe, Africa, Middle East and USA
68-73 days growing days
- Heavy, white dense heads
- Strong plant vigour
- Very good self protection

Variety	Segment	Resistances
Space Dream SGC2095	Fresh	

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

syngenta[®]

Powered by
Syngenta Vegetable Seeds

Cauliflower



Highlights
video



Spacestar Gold SGC2142

- Medium cycle variety for Africa and Middle East
- Successor of Spacestar
- 70-75 days growing days
- Strong plant vigour
- Nice curd quality
- For summer and autumn production

Variety	Segment	Resistances
Spacestar Gold SGC2142	Fresh	

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

syngenta[®]

Powered by
Syngenta Vegetable Seeds

Cauliflower



Variety
video



Wonder SGC3096

- High yield variety for late harvest 90-100 growing days
- Good performance in North and South Europe
- Fresh market and processing industry
- Very firm heads with white dense florets
- Healthy plants with very good self protection

Variety	Segment	Resistances
Wonder SGC3096	Fresh	

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

syngenta[®]

Powered by
Syngenta Vegetable Seeds

Kohlrabi

▶
Variety
video



Bogart SGK0008

- Medium maturity variety (65-70 growing days) for Europe, Africa and Middle East
- Flattish shape of the bulb
- Erect leaves and long stem support harvest
- Perfect for summer production and weaker soil

Variety	Segment	Resistances
Bogart SGK0008	General	HR:Pb

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

syngenta®

Powered by
Syngenta Vegetable Seeds

Kohlrabi

▶
Variety
video



Brando SGK0024

- Early variety for spring production in tunnel and open field for Europe, Africa and Middle East
- Erected plant suitable to high density transplanting
- Nice flat bulb
- Easy to harvest with healthy bottom

Variety	Segment	Resistances
Brando SGK0024	General	

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

syngenta®

Kohlrabi



Fonda SGK0007

- Medium maturity variety (63-68 growing days) for Europe, Africa and Middle East
- Very strong root system
- Erect leaves and long stem support fast harvest
- Late spring, autumn and summer production

Variety	Segment	Resistances
Fonda SGK0007	General	HR:Pb

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

Brussels Sprouts



Hey Melis

- Early variety with strong bolting tolerance for Europe
- Vigorous healthy plants
- High quality sprouts with dark colour
- November harvest for fresh market and processing industry

Variety	Segment	Resistances
Hey Melis	Fresh	

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

Brussels Sprouts



Marcantus

- High yielding variety for Europe
- Very good deleafing, good sturdiness
- High quality sprouts with less buds
- October/November harvest for fresh market and processing industry

Variety	Segment	Resistances
Marcantus	Fresh	

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

Powered by
Syngenta Vegetable Seeds

White Cabbage



Variety
video



Highlights
video



Eldoret SGW0431

- Medium, fresh market variety in Fabiola segment for Africa and Middle East
- Maturity around 65-70 days
- Round heads weight 3-4 kg
- Suitable for continental growing conditions
- Strong against thrips and xanthomonas

Variety	Segment	Resistances
Eldoret SGW0431	Fresh	HR: Foc 1

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

syngenta[®]

Powered by
Syngenta Vegetable Seeds

White Cabbage



Variety
video



Pruktor Cemes

- Medium, fresh market for Africa and Middle East
- Successor of Pruktor with better uniformity
- Broad adaptation to difficult growing conditions
- Healthy plants
- Head weight 3-4 kg

Variety	Segment	Resistances
Pruktor Cemes	Fresh	HR: Foc 1, Strong against Tt

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

syngenta[®]

White Cabbage



Bolikor

- Early variety for open field production for north Europe
- Around 60-65 growing days
- Nice round dark green head weight 2-2,5 kg
- Long field standing ability
- Suitable to grow in difficult early spring conditions

Variety	Segment	Resistances
Bolikor	Fresh	HR: Foc 1

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

Powered by
Syngenta Vegetable Seeds

White Cabbage



Variety
video



Rosberg SGW0541

- Adaptability to cold spring conditions tunnel and open field in north Europe
- 50-55 growing days, strong bolting tolerance
- Good field standing ability and density of heads
- Round heads weight 1,2-1,8 kg

Variety	Segment	Resistances
Rosberg SGW0541	Fresh	

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

syngenta[®]

Powered by
Syngenta Vegetable Seeds

White Cabbage



Variety
video



Successor SGW0359

- Medium fresh and short storage for north Europe
- Variety in Agressor segment, recommended for areas with higher pressure of Xanthomonas
- More round heads weight 3-5 kg, perfect for supermarkets
- Fresh market and processing
- Excellent performance in continental conditions

Variety	Segment	Resistances
Successor SGW0359	Storage	HR: Foc 1, Strong against Tt

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

syngenta[®]

Powered by
Syngenta Vegetable Seeds

White Cabbage



Variety
video



Webber SGW0508

- Most reliable for first open field production for north Europe
- 50-55 growing days
- Strong plant vigour
- Round heads weight 1,3-1,8kg
- Good field standing ability

Variety	Segment	Resistances
Webber SGW0508	Fresh	

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

syngenta[®]

White Cabbage



Katator SGW0303

- Quisor type for summer production for Europe
- Very strong against thrips
- Head weight 2,5-3,5 kg
- Fresh market variety

Variety	Segment	Resistances
Katator SGW0303	Fresh	HR: Foc 1, Strong against Tt

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

White Cabbage



Terminator SGW0247

- New medium variety for Europe in continental stress growing conditions
- Maturity 100-110 days
- Head size 4-5 kg
- Strong ,healthy plant
- Strong against thrips

Variety	Segment	Resistances
Terminator SGW0247	Fresh	HR: Foc 1, Strong against Tt

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

White Cabbage



Variety
video



Highlights
video



Toreador SGW0417

- Medium fresh segment variety for north Europe
90-95 growing days
- Head weight 2-3 kg, excellent for supermarkets
- Strong against thrips, outstanding performance across different geographies
- Upright leaves and high uniformity
- Perfect for mechanical harvest

Variety	Segment	Resistances
Toreador SGW0417	Fresh	HR: Foc 1, Strong against Tt

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

White Cabbage



Kilastress SGW0377

- Late variety for north Europe 125-130 days
- Very high yield potential, head weight 4-5 kg
- Good vigour and very long field standing ability
- Short storability, until March

Variety	Segment	Resistances
Kilastress SGW0377	Processing	HR: Pb

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

Powered by
Syngenta Vegetable Seeds

White Cabbage



Variety
video



Kepler

- Long storage variety for north Europe
- Big heads 3-4 kg
- Keeps green color after storage
- Broad adaptability to growing conditions

Variety	Segment	Resistances
Kepler	Storage	HR:foc1

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

syngenta[®]

White Cabbage



Variety
video



Highlights
video



Storidor

- Late variety for long storage for Europe
- 120-125 growing days
- Marketable heads can reach 3-4 kg
- Erect plant, strong root system and high uniformity makes variety suitable to mechanical harvest

Variety	Segment	Resistances
Storidor	Storage	HR: Foc 1

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

Red Cabbage



Red River SGX0057

- Dark red heads with very good internal structure
- Fresh market variety for winter production in South Europe and long storage in North Europe
- High round, waxy heads weight 2-3 kg)
- Very good standing ability

Variety	Segment	Resistances
Red River SGX0057		HR: Foc 1, IR: Mb

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

Powered by
Syngenta Vegetable Seeds

Savoy Cabbage



Variety
video



Atractum SGS0325

- Fast variety with high yield (60-65 days) for north Europe
- Head weight 1-2,5 kg depending on density
- Very good field standing ability - strong against cracking
- Spring and summer production

Variety	Segment	Resistances
Atractum SGS0325		IR:Ac

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

syngenta[®]

Powered by
Syngenta Vegetable Seeds

Savoy Cabbage



Coradi SGS0323

- Spring ,summer variety for fresh market for Europe
- Very good savoynes and sanitary of crop
- Full package of fungi diseases
- Good field standing ability
- Round compact heads

Variety	Segment	Resistances
Coradi SGS0323		HR:Pb,IR:Ac,;IR:Mb

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

syngenta[®]

Powered by
Syngenta Vegetable Seeds

Savoy Cabbage



Cordesa

- Summer variety for fresh market in Europe
- Broad package of resistances
- Easy to pack
- Heavy dense heads

Variety	Segment	Resistances
Cordesa		HR:Pb,IR:Ac,:IR:Mb

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

syngenta[®]

Powered by
Syngenta Vegetable Seeds

Savoy Cabbage



Corripa SGS0294

- Very late variety for fresh market for Europe
- Broad package of resistances
- Frost tolerance
- Dark green heads weight 1-1,5 kg

Variety	Segment	Resistances
Corripa SGS0294		HR:Pb,IR:Ac,:IR:Mb

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

syngenta[®]

Powered by
Syngenta Vegetable Seeds

Savoy Cabbage



Variety
video



Milandina SGS0318

- Summer type (80-90 days) for Europe
- Oval heads 1,5-2,0kg
- Good savoyness
- Very good field standing ability

Variety	Segment	Resistances
Milandina SGS0318		IR:Ac

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

syngenta[®]

Savoy Cabbage



Manitoba

- Very late variety for north Europe
- Season extension for Kamchatka
- Round heads weight 1-1,5 kg
- Very healthy plant until end of season
- Strong frost and bolting tolerance

Variety	Segment	Resistances
Manitoba		

**For more information and specific details,
please contact your local Technical Sales Representative.**


Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation



**You can always
visit us again
anytime you want**

foi.syngentavegetables.com

syngenta®