

Fields of Innovation

September 21 - 25, 2020

Event

Powered by Syngenta
Vegetable Seeds

Thank you
for visiting
and enjoy your
Field Book!

syngenta®

Content of your Field Book

EL Bravo LDSP999

EL Furio LDSP996

Odysseus LDSP948

Annisole LS17703

Coriole LS16743

Frisole LS16744

Laguiole LS17701

Tirole LOBA19-0018

Elodian LS17650

Radian

Canagio LS18468

Prodigio LS18467

Rubagio LS18469

Primice LMUL18-0010

Ice Circle LS15542



Content of your Field Book

Ice Desert LS16528

Ice Jazz LS17504

Ice Palace

Ice Party LS17510

Ice Wave LS15548

Icebazaar PYB7101A

Actina LS9858

Belendra LS16820

Calorina LS16847

Casual LS17833

LRMN19-1959 LRMN19-1959

Typical LS16841

Presidential LS8832

Annita LS16042

Claudita LS15028



Content of your Field Book

Crispita II LS15012

Cupidita LS18061

Icevita LS17053

LBAB19-1938 LBAB19-1938

Red Crispita II LBAB18-0020

LMUL18-0013 LMUL18-0013

LMUL19-1922 LMUL19-1922



Spinach



EL Bravo LDSP999

- Good variety for all of Europe
- Thick leaves and dark green, fresh green colour
- Good speed growth for spring and autumn period
- Variety close to EL Tajin with better Peronospora resistance

Variety	Segment	Resistances
EL Bravo LDSP999	Babyleaf	Pfs: 1-7,9-17 + Sb (HR)

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

Powered by
Syngenta Vegetable Seeds

Spinach


Variety
video



EL Furio LDSP996

- Our new candidate for babyleaf production with dark green colour, slow growth speed and thick leaf
- Good growth speed (not too fast) with good harvesting window
- Very nice dark green, fresh color with nice leaf shape and good 3D

Variety	Segment	Resistances
EL Furio LDSP996	Babyleaf	Pfs: 1-7,9-17 + Sb (HR)

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation



Spinach



Odysseus LDSP948

- Robust and flexible spinach variety for all markets : babyleaf, bunching and also processing
- High yield due to thick leaves
- Round to oval leaf shape with good leaf colour
- Good bolting tolerance for processing market

Variety	Segment	Resistances
Odysseus LDSP948	Babyleaf	Pfs: 1-9,11-16 + Sb (HR)

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

Powered by
Syngenta Vegetable Seeds

Batavia



Annisole LS17703

- For all of Europe (North & South)
- Very attractive colour (glossy) and nice upside presentation (ondulée leaf type)
- Uniform and flexible variety
- Semi erect plant and healthy bottom with small core => easy to harvest
- Large harvesting window

Variety	Segment	Resistances
Annisole LS17703	General	Bl:16-35EU, Nr:0 (HR)

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

syngenta®

Batavia



Coriole LS16743

- Dark green Batavia with vibrant colour
- For fresh and processing, North Europe (late spring and autumn), South East Europe (spring, autumn and winter)
- High yield potential
- Perfect in combination with Parole to supply all season processing

Variety	Segment	Resistances
Coriole LS16743	General	Bl:16-35EU, Nr:0 (HR) - LMV:1(IR)

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

Powered by
Syngenta Vegetable Seeds

Batavia



Frisole LS16744

- Curly type with very nice color and upside presentation for fresh market
- Very uniform variety with good plant organisation
- Easy to harvest due to semi erect habit and healthy bottom

Variety	Segment	Resistances
Frisole LS16744	General	Bl:16-35EU, Nr:0 (HR) - LMV:1(IR)

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

syngenta®

Batavia



Laguiole LS17701

- For fresh market (outdoor), North West Europe (summer), South East Europe (late spring, summer, early autumn)
- Well organized plant and very good presentation for fresh market
- Easy to harvest : semi erect plant, healthy bottom and small core
- Very uniform variety

Variety	Segment	Resistances
Laguiole LS17701	General	Bl:16-35EU, Nr:0 (HR) - LMV:1(IR)

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

Powered by
Syngenta Vegetable Seeds

Batavia



Tirole LOBA19-0018

- Good vigour for all of Europe, Spring and Autumn period with nice, fresh green color
- Good upside presentation with good heart filling
- Good and semi erect bottom quality : makes it easy to harvest

Variety	Segment	Resistances
Tirole LOBA19-0018	General	BI:16-36EU, Nr:0, Fol:1 (HR) - LMV:1 (IR)

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

syngenta[®]

Butterhead



Elodian LS17650

- Full mildew and Nr:0 resistances
- Very flexible variety for almost all season (outdoor - north west of Europe)
- Smaller to medium size for fresh market, perfect for single packs
- Good amount of frame leaves and easy to cut and clean => security against soil borne diseases

Variety	Segment	Resistances
Elodian LS17650	General	Bl:16-36EU, Nr:0, TBSV (HR) - LMV:1(IR)

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

Butterhead



Radian

- Full mildew package and Nr:0 resistances
- For fresh and processing market, all of Europe (summer)
- Very good upside with attractive green color and nice / healthy bottom quality and good behaviour against internal tipburn and bolting (real summer variety)
- Good filling offering high white rate (for processing market) and good harvesting window

Variety	Segment	Resistances
Radian	General	Bl:16-36EU, Nr:0, TBSV (HR) - LMV:1(IR)

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

Bellagio



Canagio LS18468

- Green butterhead Bellagio variety for North West Europe, spring -> fall
- Full breimia package and nasonovia resistance
- Good volume and yield for processing market with nice green leaf colour and shape
- Flexible / rustic variety that works well in different season and locations (outdoor production)

Variety	Segment	Resistances
Canagio LS18468	General	BI: 16-36EU, Nr: 0 (HR)

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

Powered by
Syngenta Vegetable Seeds

Bellagio



Variety
video



Prodigio LS18467

- Good red, incised variety with full Bremia resistance package and Nasonovia Nr:0 resistance
- Red bellagio with good plant organisation combining nice red and green leaf colour, good leaf shape, branching and yield potential
- Rustic and flexible variety for all EU

Variety	Segment	Resistances
Prodigio LS18467	General	BI: 16-36EU, Nr: 0 (HR)

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

syngenta[®]

Bellagio



Rubagio LS18469

- Red butterhead Bellagio variety for North West Europe, spring -> autumn
- Full breimia package and nasonovia resistance
- Very nice and glossy red leaf colour and shape
- To be tested during 2020 and 2021 season in key markets (Europe and Australia)

Variety	Segment	Resistances
Rubagio LS18469	General	BI: 16-36EU, Nr: 0 (HR)

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

Powered by
Syngenta Vegetable Seeds

Bellagio



Primice LMUL18-0010

- New green incised variety for Hydroponic production for North-West Europe (Nordics) and Japan
- For spring, autumn and winter production
- Very robust and flexible variety (with good bolting, tipburn and twisting behaviour)
- Works well with 1 seed / pot but also with 2 or 3 seeds / pot as well

Variety	Segment	Resistances
Primice LMUL18-0010	Krispice	Bl:16-36EU, TBSV (HR) - LMV:1(IR)

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

syngenta[®]

Powered by
Syngenta Vegetable Seeds

Iceberg

▶
Variety
video



Ice Circle LS15542

- Uniform head shape and concentrated maturity for highest harvesting rate
- Quick filling head which becomes not too dense/hard thanks very regular layers
- Very well organized plant with good internal filling
- Reliable and rustic variety for all of Europe, Spring and Autumn period

Variety	Segment	Resistances
Ice Circle LS15542	General	Bl:16-36EU, Nr:0, Fol:4, TBSV (HR)

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

syngenta®

Powered by
Syngenta Vegetable Seeds

Iceberg



Ice Desert LS16528

- Fresh market for Africa and Middle-East
- New summer variety to support Icebazaar in Summer period
- Good Glossy color and head shape
- To be tested in 2020 & 2021 locally

Variety	Segment	Resistances
Ice Desert LS16528	General	

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

syngenta®

Powered by
Syngenta Vegetable Seeds

Iceberg



Ice Jazz LS17504

- For central Europe and Turkey
- Autumn variety with full mildew and Nasonovia resistance
- Very good bottom quality (smooth nerves) and nice head shape & filling

Variety	Segment	Resistances
Ice Jazz LS17504	General	Bl:16-36EU, Nr:0 (HR)

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

syngenta®

Iceberg



Ice Palace

- Salinas iceberg type recommended for autumn and spring harvest
- Excellent head shape and size / calibre.
- Very good bottom quality to avoid filming issues and browning later on
- Good tip burn and bolting tolerance

Variety	Segment	Resistances
Ice Palace	General	BI: 16-29, 32, 34, 36EU, Nr: 0, Fol: 1, TBSV (HR)

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

Powered by
Syngenta Vegetable Seeds

Iceberg

Variety
video



Ice Party LS17510

- For central Europe (spring & autumn), north Europe (almost all year)
- A new iceberg variety with full mildew and Nasonovia resistance
- Bigger head size for fresh and processing markets
- Really long harvesting window
- Good shape for easy picking
- Good complementarity with Ice Circle and Ice Music

Variety	Segment	Resistances
Ice Party LS17510	General	Bl:16-36EU, Nr:0, TBSV (HR)

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

syngenta[®]

Powered by
Syngenta Vegetable Seeds

Iceberg



Ice Wave LS15548

- Robust and reliable iceberg with full mildew resistance
- Good bolting and tip burn tolerance (also in difficult conditions)
- Semi-flat head shape and flat base for easy picking and packing
- Large planting and harvesting window Spring - Autumn (North West of EU : almost all the season)
- Reliable and flexible variety

Variety	Segment	Resistances
Ice Wave LS15548	General	BL:16-36EU, Nr:0, TBSV (HR)

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

syngenta®

Powered by
Syngenta Vegetable Seeds

Iceberg



Icebazaar PYB7101A

- Our best summer iceberg variety for Africa and Middle East area
- Very good heat tolerance (with good bolting and tipburn behaviour)
- Good growth speed and a large harvesting window
- Head size and shape is perfect for fresh market

Variety	Segment	Resistances
Icebazaar PYB7101A	General	BI:17EU, TBSV (HR)

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

syngenta®

Powered by
Syngenta Vegetable Seeds

Romaine



Variety
video



Actina LS9858

- Setting the highest standard for tip burn tolerance in Midi romaine market in Europe
- Very reliable production all outdoor season
- Long harvesting window and suitable for both markets, fresh and processing

Variety	Segment	Resistances
Actina LS9858	Midi	Bl:16-28,30-32EU, Nr:0, Fol:1, Fol:4 (HR)

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

syngenta[®]

Romaine



Belendra LS16820

- Intensive red colour leaf and good red colour penetration
Well shaped romaine between gem and mini types (Spain)
- Reliable variety with consistent results and easy to cut
- Good head filling
- For central Europe (spring and autumn), south Europe for spring and autumn as mini Romain and winter as Red Gem

Variety	Segment	Resistances
Belendra LS16820	Mini	BI:16-35EU (HR)

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

Romaine



Calorina LS16847

- Variety with good bolting tolerance for warm conditions
- Good plant organisation and good bottom quality
- Variety with good closing head and with good heart filling (for processing market)
- Variety with healthy bottom: easy to harvest
- Potential for all of Europe, north to be tested in 2020/2021

Variety	Segment	Resistances
Calorina LS16847	Midi	BI:16-36EU (HR)

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

Powered by
Syngenta Vegetable Seeds

Romaine



Casual LS17833

- Our new Midi romaine variety with full breimia package for North and South of Europe (EU countries)
- Good head shape with good filling
- Good and healthy bottom quality => easy to cut

Variety	Segment	Resistances
Casual LS17833	Midi	Bl:16-36EU, Nr:0, TBSV (HR)

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

syngenta®

Powered by
Syngenta Vegetable Seeds

Romaine



LRMN19-1959

- Mini red romaine with excellent colour
- Suitable for all season production in the UK
- Nice straight shape and easy to harvest”

Variety	Segment	Resistances
LRMN19-1959	Mini	BI:16-36EU (HR)

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

syngenta®

Romaine



Typical LS16841

- Taller Midi romaine compare to Actina with high flexibility for mainly Spring and Autumn production
- Better bolting tolerance than Actina
- Darker colour and perfect plant shape (with good behaviour against twisting)

Variety	Segment	Resistances
Typical LS16841	Midi	Bl:16-36EU, Nr:0, Fol:1, TBSV (HR)

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

Powered by
Syngenta Vegetable Seeds

Romaine



Variety
video



Presidential LS8832

- Our unic romaine variety for Turkey market
- Presidential is working all year round in open field & also passive green house
- Presidential have a very nice upside presentation and is a rustic variety to garante your production

Variety	Segment	Resistances
Presidential LS8832	Midi	Bl: 16-28,30-32EU, Fo: 1 (HR)

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

syngenta[®]

Babyleaf



Annita LS16042

- Slow growing batavia with attractive and intensive green colour
- Better harvesting window than Lucita and Sensita (slow speed growth)
- Thick and flexible leaves offering high yield potential
- Nice green color and good leaf shape / 3D

Variety	Segment	Resistances
Annita LS16042	Babyleaf	BI:16-36EU (HR)

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

Babyleaf



Claudita LS15028

- Deep red glossy Batavia with good 3D shape
- Narrow stem that avoids oxidation on cutting point
- Full Bremia and Fusarium high resistance
- To support Izabita and Lennita (if needed)

Variety	Segment	Resistances
Claudita LS15028	Babyleaf	Bl:16-36EU, Fol:1, Fol:4 (HR)

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

Babyleaf



Variety
video



Crispita II LS15012

- Unique leaf shape and very good 3D structure with crispy texture
- Good leaf shape and color with thick leaf - helps to maintain plastic bag volume and provide long shelf life
- Good yield, excellent taste and wide harvesting window

Variety	Segment	Resistances
Crispita II LS15012	Babyleaf	BI:16-36EU, Fol:4 (HR) - LMV:1(IR)

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

Babyleaf



Variety
video



Cupidita LS18061

- Our new red romaine variety : our aim is to develop a more vigorous variety compare to Kerrita
- Cupidita is slightly faster than Kerrita but it keeps the same nice red and glossy color, the same leaf shape and leaf thicknes
- It is an improvement on Kerrita during cold conditions due to its faster growth speed

Variety	Segment	Resistances
Cupidita LS18061	Babyleaf	Bl:16,18-36EU, Fol:1 (HR)

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

Babyleaf



Variety
video



Icevita LS17053

- Our new variety in Baby Iceberg type - innovative leaf typology
- Leaves thickness is offering crunchiness (close to Iceberg type)
- Leaves also have a good 3D shape to offer good volumes in the bag

Variety	Segment	Resistances
Icevita LS17053	Crispy	Bl:16-36EU, Nr:0, TBSV (HR)

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

Babyleaf



LBAB19-1938

- Our new variety in Green Crispy type in order to support Crispita II, Crackita and Mesclita varieties
- Very nice and attractive green color with good leaf shape and very good 3D
- Thick leaves and good growth speed (not too fast) to secure yield and correct harvesting window

Variety	Segment	Resistances
LBAB19-1938	Crispy	Bl:16-36EU, TBSV (HR)

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

Babyleaf



Variety
video



Red Crispita II LBAB18-0020

- Or new Red Crispita II to replace initial red crispita
Good complementarity with Crispita II and Crackita
- Thick leaves that provide good yield and good 3D in packaging
(behave like back bone in the plastic bag)
- Good shelf life and good leaf texture (crunchiness) thanks to
thicker leaves
- Very good harvesting window

Variety	Segment	Resistances
Red Crispita II LBAB18-0020	Babyleaf	Bl:16-36EU, TBSV (HR)

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

Powered by
Syngenta Vegetable Seeds

Bellagio



LMUL18-0013

- New red bellagio (red incised type), in all of Europe for summer
- Nice red leaf color, good branching and incised leaf
- Premium product for summer segment for UK, Australia and US market to be tested during 2020 / 2021 season locally

Variety	Segment	Resistances
LMUL18-0013	General	BI: 16-36EU, Nr: 0 (HR)

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

syngenta®

Batavia



Variety
video



Highlights
video



LMUL19-1922

- Our new, innovative type targeting mainly burgers
- Variety offering a high % of usable leaves. Leaves fit well in a burger (round shape and same leaf size)
- Leaves with good taste and texture (crunchiness)
- Leaves also offering good heat tolerance (to keep leaf texture and crunchiness in the burger when in contact with hot food)

Variety	Segment	Resistances
LMUL19-1922	Crispy	BI:16,18-21,23-26,28-36EU (HR)

**For more information and specific details,
please contact your local Technical Sales Representative.**

Syngenta Seeds Vegetables has exercised reasonable care and skill in compiling this brochure. All resistances quoted refer only to strains of races or pathotypes indicated on the varieties. Other pathogen races or pest biotypes capable of overcoming the resistance may exist or emerge. The Syngenta resistance against Club Root is effective against the predominant races Pb:0 and Pb:1 and against the less frequent race Pb:3 but not against the infrequent race Pb:2 that may occur in some fields. Genetic resistance is only one of the tools to manage Club Root. Culture measures such as liming, use of fertilizers with high percentage of calcium, proper drainage, good crop hygiene management are several of important components of an integrated approach to manage the disease. Syngenta Seeds Vegetables uses established analytical methods to verify specific variety resistances. However, host specificity of pests or pathogens may vary depending on environmental factors. In order to maximize the efficiency of a resistance, it is highly recommended to combine different ways of control such as growing conditions, plant protection products and genetic resistance as part of an integrated crop management. All data in this brochure are intended for general guidance only and the user should apply it in accordance with his own knowledge and experience of local conditions. In case of doubt we recommend that a small scale trial production be carried out to determine how local conditions may affect the variety.

Syngenta Seeds Vegetables cannot accept any liability in connection with this brochure.

syngentavegetables.com



#fieldsofinnovation

syngenta[®]



**You can always
visit us again
anytime you want**

foi.syngentavegetables.com

syngenta[®]